A Minimal Type II Seesaw Model

Pei-Hong Gu,* He Zhang,† and Shun Zhou‡
Institute of High Energy Physics, Chinese Academy of Sciences, P.O. Box 918 (4), Beijing 100049, China§

We propose a minimal type II seesaw model by introducing only one right-handed neutrino besides the $SU(2)_L$ triplet Higgs to the standard model. In the usual type II seesaw models with several right-handed neutrinos, the contributions of the right-handed neutrinos and the triplet Higgs to the CP asymmetry, which stems from the decay of the lightest right-handed neutrino, are proportional to their respective contributions to the light neutrino mass matrix. However, in our minimal type II seesaw model, this CP asymmetry is just given by the one-loop vertex correction involving the triplet Higgs, even though the contribution of the triplet Higgs does not dominate the light neutrino masses. For illustration, the Fritzsch-type lepton mass matrices are considered.

PACS numbers: 11.30.Fs, 14.60.Pq, 14.60.St, 14.80.Cp

arXiv:hep-ph/0606302v3 29 Sep 2006

Recent neutrino oscillation experiments have provided us with very convincing evidence that neutrinos are massive and lepton flavors are mixed [1]. In the three-neutrino mixing scheme, a global analysis of experimental data yields: two independent neutrino mass-squared differences $\Delta m^2_{21} = (7.2 \sim 8.9) \times 10^{-5}$ eV^2, $|\Delta m^2_{32}| = (1.7 \sim 3.3) \times 10^{-3}$ eV^2 and three mixing angles $30^\circ < \theta_{12} < 38^\circ$, $36^\circ < \theta_{23} < 54^\circ$, $\theta_{13} < 10^\circ$ at the 99% confidence level [2].

On the other hand, the cosmological baryon asymmetry, which is characterized by the ratio of baryon to photon number densities, has also been measured by the WMAP experiment to a very good precision [3]

$$\eta_{\rm B} \equiv \frac{n_{\rm B}}{n_\gamma} = (6.1 \pm 0.2) \times 10^{-10} \; . \tag{1}$$

Sometimes the baryon asymmetry is also represented by $Y_{\rm B} \equiv n_{\rm B}/s = \eta_{\rm B}/7.04 \simeq (8.4 \sim 8.9) \times 10^{-11}$ with s being the entropy density.

Leptogenesis [4] is now an attractive scenario to simultaneously explain the neutrino oscillation phenomena and the baryon asymmetry through the famous seesaw mechanism [5]. In the ordinary type I seesaw models, the leptogenesis scenario can be realized by introducing two or more right-handed neutrinos with heavy Majorana masses to the $SU(2)_L \times U(1)_Y$ standard model. Another interesting alternative is to consider the so-called type II seesaw models [6], in which the $SU(2)_L$ triplet Higgs, besides the right-handed neutrinos, can also contribute to the light neutrino masses as well as the baryon asymmetry generation.

In this paper, we propose a minimal type II seesaw model by extending the standard model with only one right-handed neutrino in addition to the $SU(2)_L$ triplet Higgs.[1] The CP asymmetry in the decay of the right-handed neutrino just arises from the interference between the tree level diagram and the one-loop vertex correction involving the triplet Higgs. At the same time, the light neutrino mass matrix can receive comparable contributions from the triplet Higgs and the right-handed neutrino. For illustration, we will consider Fritzsch-type lepton mass matrices with five texture zeroes in the specific discussions and calculations. It will be shown that our model can simultaneously explain the neutrino properties and the baryon asymmetry.

It is convenient to start with our discussions from the general type II seesaw models [7, 8]:

$$\begin{aligned} -\mathcal{L} &= M^2_\Delta {\rm Tr}\Delta^\dagger_L \Delta_L + g_{\alpha\beta}\bar{\psi}^C_{L\alpha} i\tau_2 \Delta_L \psi_{L\beta} - \mu \phi^T i\tau_2 \Delta_L \phi \\ &\quad + \frac{1}{2} M_i \bar{\nu}^C_{Ri} \nu_{Ri} + y_{\alpha i} \bar{\psi}_{L\alpha} \nu_{Ri} \phi + h.c. \\ &= M^2_\Delta {\rm Tr}\Delta^\dagger_L \Delta_L + \frac{1}{2} M_i \bar{N}_i N_i + g_{\alpha\beta}\bar{\psi}^C_{L\alpha} i\tau_2 \Delta_L \psi_{L\beta} \\ &\quad - \mu \phi^T i\tau_2 \Delta_L \phi + y_{\alpha i} \bar{\psi}_{L\alpha} N_i \phi + {\rm h.c.} \, . \end{aligned} \tag{2}$$

Here $\psi_\alpha = (\nu_\alpha, l_\alpha)^T$ $(\alpha = e, \mu, \tau)$, $\phi = (\phi^0, \phi^-)^T$ are the lepton and the Higgs doublets, while

$$\Delta_L = \begin{pmatrix} \frac{1}{\sqrt{2}}\delta^+ & \delta^{++} \\ \delta^0 & -\frac{1}{\sqrt{2}}\delta^+ \end{pmatrix}$$

is the triplet Higgs. ν_{Ri} $(i = 1, ..., d)$ are the right-handed neutrinos, and $N_i = \nu_{Ri} + \nu^C_{Ri}$ is defined as the heavy Majorana neutrinos. We have conveniently chosen the basis, in which the Majorana mass matrix of right-handed neutrinos is diagonal, *i.e.* $M = {\rm Diag}\{M_1, ..., M_d\}$. Obviously, g is generally a complex 3×3 matrix. In the usual type II seesaw models, there are several right-handed neutrinos, *i.e.* $d \geq 2$. However, only one right-handed neutrino and hence three Yukawa couplings $y_{\alpha 1}$ $(\alpha = e, \mu, \tau)$ occur in our minimal type II seesaw model with $d = 1$.

*Electronic address: guph@mail.ihep.ac.cn
†Electronic address: zhanghe@mail.ihep.ac.cn
‡Electronic address: zhoush@mail.ihep.ac.cn
§Mailing address

[1] Here the *minimal* is in the sense of the number of new particles compared with the conventional type II seesaw model.



FIG. 1: The one-loop diagrams of N_i decay.

After the electroweak phase transition, the mass matrix of the light neutrinos $\nu_\alpha = \nu_{L\alpha} + \nu^C_{L\alpha}$ can be written as

$$M_\nu = -y^* \frac{1}{M} y^\dagger v^2 + 2 g v_L = M_\nu^{\rm I} + M_\nu^{\rm II} \,, \tag{3}$$

where $M_\nu^{\rm I}$ is the ordinary type I seesaw mass term, $M_\nu^{\rm II}$ is the type II seesaw mass term. $v = 174\,{\rm GeV}$, $v_L \simeq \mu^* v^2/M_\Delta^2$ are the vacuum expectation values of ϕ and Δ_L, respectively. It is easy to see that v_L is naturally seesaw suppressed if Δ_L is very heavy.

The CP asymmetry ε_{N_i} from the decay of the heavy Majorana neutrinos N_i is given by the interference of the ordinary tree level decay with the three diagrams of Fig. 1. The first two diagrams are the self-energy and vertex correction involving the heavy Majorana neutrinos, while the third is due to the contribution from the triplet Higgs. Under the assumption that $M_1 \ll M_2, ..., M_d, M_\Delta$, the CP asymmetry ε_{N_1} can be simplified as [8, 9]:

$$\varepsilon_{N_1} = \varepsilon^N_{N_1} + \varepsilon^\Delta_{N_1} \,, \tag{4}$$

$$\varepsilon^N_{N_1} \simeq \frac{3}{16\pi}\frac{M_1}{v^2}\frac{\sum_{\alpha\beta} {\rm Im}\left[y^{\nu\dagger}_{1\alpha} y^{\nu\dagger}_{1\beta} \left(M_\nu^{\rm I*}\right)_{\alpha\beta}\right]}{\left(y^{\nu\dagger} y^\nu\right)_{11}} \,, \tag{5}$$

$$\varepsilon^\Delta_{N_1} \simeq \frac{3}{16\pi}\frac{M_1}{v^2}\frac{\sum_{\alpha\beta} {\rm Im}\left[y^{\nu\dagger}_{1\alpha} y^{\nu\dagger}_{1\beta} \left(M_\nu^{\rm II*}\right)_{\alpha\beta}\right]}{\left(y^{\nu\dagger} y^\nu\right)_{11}} \,. \tag{6}$$

It is shown that the contributions of N_i (diagrams (a) and (b) in Fig. 1) and Δ_L (diagram (c) in Fig. 1) to ε_{N_1} are proportional to their respective contributions to the light neutrino mass matrix [8]. Therefore, the contribution from the triplet Higgs to the above CP asymmetry can be neglected in the limit $M_\nu^{\rm II} \ll M_\nu^{\rm I}$. However, this analysis is invalid in our minimal type II seesaw model, since there is just one heavy Majorana neutrino, the one-loop vertex correction involving the triplet Higgs is the sole source of this CP asymmetry and we always obtain

$$\varepsilon_{N_1} \equiv \varepsilon^\Delta_{N_1} \,, \tag{7}$$

even though the contribution of Δ_L does not dominate the light neutrino mass.

For illustration, we simply assume the Fritzsch-type textures [10] of $M_\nu^{\rm II}$ and the charged lepton mass matrix M_l:

$$M_\nu^{\rm II} = v_L \begin{pmatrix} 0 & C_\nu e^{i\alpha_\nu} & 0 \\ C_\nu e^{i\alpha_\nu} & 0 & B_\nu e^{i\beta_\nu} \\ 0 & B_\nu e^{i\beta_\nu} & A_\nu e^{i\gamma_\nu} \end{pmatrix} \tag{8}$$

and

$$M_l = v \begin{pmatrix} 0 & C_l e^{i\alpha_l} & 0 \\ C_l e^{i\alpha_l} & 0 & B_l e^{i\beta_l} \\ 0 & B_l e^{i\beta_l} & A_l e^{i\gamma_l} \end{pmatrix} \,, \tag{9}$$

where $A_{\nu,l}$, $B_{\nu,l}$ and $C_{\nu,l}$ are real and positive. In addition, we set

$$y = i y_0 \left(0, r, 1\right)^T \,, \tag{10}$$

where the imaginary unit i has been inserted to cancel the minus sign in front of the type I term for convenience. Substituting Eqs. (8) and (10) into Eq. (3), we can obtain the effective neutrino mass matrix

$$M_\nu = m_0 \begin{pmatrix} 0 & \widehat{C} e^{i\alpha_\nu} & 0 \\ \widehat{C} e^{i\alpha_\nu} & r^2 & r + \widehat{B} e^{i\beta_\nu} \\ 0 & r + \widehat{B} e^{i\beta_\nu} & 1 + \widehat{A} e^{i\gamma_\nu} \end{pmatrix} \,, \tag{11}$$

where $m_0 \equiv v^2 y_0^2/M_1$ and $\widehat{A} \equiv v_L A_\nu/m_0$, likewise for $\widehat{B}$ and $\widehat{C}$. The strategies to diagonalize lepton mass matrices of the form in Eqs. (9) and (11) can be found in Refs. [10] and [11]. For simplicity, we adopt two assumptions:

$r = \sqrt{m_2/m_0}$ and $\arg\left(1+\widehat{A}e^{i\gamma_\nu}\right) = 2\arg\left(r+\widehat{B}e^{i\beta_\nu}\right)$, and then obtain

$$\begin{aligned}
\widehat{A} &= \left[\frac{(m_3-m_1)^2}{m_0^2} - \sin^2\gamma_\nu\right]^{1/2} - \cos\gamma_\nu \, , \\
\widehat{B} &= \left[\frac{m_1 m_3 (m_3-m_1-m_2)}{m_0^2(m_3-m_1)} - r^2\sin^2\beta_\nu\right]^{1/2} \\
&\quad -r\cos\beta_\nu \, , \\
\widehat{C} &= \left[\frac{m_1 m_2 m_3}{m_0^2(m_3-m_1)}\right]^{1/2} \qquad (12)
\end{aligned}$$

and

$$\begin{aligned}
A_l &= (m_\tau - m_\mu + m_e) \, , \\
B_l &= \left[\frac{(m_\mu - m_e)(m_\tau - m_\mu)(m_e+m_\tau)}{(m_\tau - m_\mu + m_e)}\right]^{1/2} \, , \\
C_l &= \left[\frac{m_e m_\mu m_\tau}{(m_\tau - m_\mu + m_e)}\right]^{1/2} \, . \qquad (13)
\end{aligned}$$

The Maki-Nakagawa-Sakata (MNS) mixing matrix is $V = U_l^\dagger U_\nu$, where $U_l^\dagger$ and U_ν are the unitary matrices used to diagonalize the lepton mass matrices: $U_l^\dagger M_l U_l^* = \mathrm{Diag}\left\{m_e, m_\mu, m_\tau\right\}$ and $U_\nu^\dagger M_\nu U_\nu^* = \mathrm{Diag}\left\{m_1, m_2, m_3\right\}$. Taking the neutrino oscillation experimental data at the 99% confidence level as input[2], the allowed regions of three mixing angles and Jarlskog invariant $\mathcal{J} \equiv \mathrm{Im}\left(V_{11}V_{22}V_{12}^*V_{21}^*\right)$ in our model are shown in Figs. 2 and 3. The phase parameters, exactly combinations of $\alpha_{l,\nu}, \beta_{l,\nu}$ and $\gamma_{l,\nu}$, are all varying in the range $[0, 2\pi)$. From Fig. 3, we can see that the smallest mixing angle θ_{13} is larger than 1° and the maximal value of CP-violation parameter $\mathcal{J}$ can reach 2.5%, which are to be strictly tested in the future neutrino oscillation experiments. Here we remark that the Fritzsch-type lepton mass matrices can serve as a good example to illustrate the features of the minimal type II seesaw model. For more phenomenological discussions about lepton mass matrices in the type-II seesaw models, see [12].

We now calculate the produced baryon asymmetry in the minimal type II seesaw model. Assuming the heavy Majorana neutrino N_1 is much lighter than the triplet Higgs Δ_L, the final lepton asymmetry, which is partially converted to the baryon asymmetry via the sphaleron process [14], will mainly come from the decay of N_1. We



FIG. 2: Allowed region of solar and atmospheric mixing angles, namely θ_{12} and θ_{23} in the standard parametrization [13].



FIG. 3: Allowed region of the mixing angle θ_{13} and Jarlskog invariant $\mathcal{J}$.

[2] For simplicity, we assume the normal mass hierarchy of light neutrinos in our numerical analysis.



FIG. 4: The solutions of the Boltzmann equations. $Y_{N_1}^{(\rm eq)}$ is the (equilibrium) number of heavy Majorana neutrino, while $Y_B = 28/79\, Y_{B-L}$ denotes the baryon. In the numerical calculations, we have taken $M_1 = 4 \times 10^{11}\,{\rm GeV}$, $m_1 = 8.6 \times 10^{-3}$ eV, $\Delta m^2_{21} = 8.3 \times 10^{-5}$ eV^2, $\Delta m^2_{32} = 2.3 \times 10^{-3}$ eV^2, $y_0 = 0.02$, $r = 0.64$ and $(\beta_\nu, \gamma_\nu) = (-100^\circ, -36^\circ)$. The predicted value for the final baryon asymmetry is $Y_B \simeq 8.5 \times 10^{-11}$.

can write down the relevant Boltzmann equations:

$$\frac{dY_{N_1}}{dz} = -\frac{z}{sH(M_1)}\left[\gamma_D + 2\gamma_{\phi,s} + 4\gamma_{\phi,t}\right] \times \left(\frac{Y_{N_1}}{Y_{N_1}^{\rm eq}} - 1\right) , \quad (14)$$

$$\frac{dY_{B-L}}{dz} = -\frac{z}{sH(M_1)}\left[\varepsilon_{N_1}\gamma_D\left(\frac{Y_{N_1}}{Y_{N_1}^{\rm eq}} - 1\right) + \frac{Y_{B-L}}{Y_l^{\rm eq}} \times \left(\frac{1}{2}\gamma_D + 2\gamma_N + 2\gamma_{N,t} + 2\gamma_{\phi,t} + \frac{Y_{N_1}}{Y_{N_1}^{\rm eq}}\gamma_{\phi,s}\right)\right] , \quad (15)$$

where $z = M_1/T$, $H(M_1)$ is the Hubble parameter at $T = M_1$, $Y_X^{(\rm eq)}$ is defined as $Y_X^{(\rm eq)} = n_X^{(\rm eq)}/s$ with $n_X^{(\rm eq)}$, s being the (equilibrium) number density and the entropy density, respectively. γ_X, which has been calculated by many authors [15], is the reaction density of the relevant process.

As pointed out in the foregoing discussions, because there are no other heavy Majorana neutrinos $N_i\,(i \neq 1)$, the CP asymmetry ε_{N_1} just comes from the interference between the tree level diagram and the vertex correction induced by the triplet Higgs Δ_L. For $M_\Delta \gg M_1$, we obtain:

$$\varepsilon_{N_1} \equiv \varepsilon^{\Delta}_{N_1} \simeq \frac{3}{16\pi}\frac{M_1}{v^2}\frac{\sum_{\alpha\beta}{\rm Im}\left[y^{\nu\dagger}_{1\alpha}y^{\nu\dagger}_{1\beta}\left(M_\nu^{\rm II*}\right)_{\alpha\beta}\right]}{\left(y^{\nu\dagger}y^{\nu}\right)_{11}} = \frac{3}{16\pi}\frac{y_0^2}{1+r^2}\left[\widehat{A}\sin\gamma_\nu + 2\widehat{B}r^2\sin\beta_\nu\right] . \quad (16)$$

In the last step, Eqs. (8) and (10) have been used.

Given appropriate parameters consistent with the neutrino oscillations, both the CP asymmetry ε_{N_1} and the reaction density γ_X can be computed, and then the Boltzmann equations can also be solved. For example, we take $M_1 = 4 \times 10^{11}\,{\rm GeV}$, $\Delta m^2_{21} = 8.3 \times 10^{-5}$ eV^2, $\Delta m^2_{32} = 2.3 \times 10^{-3}$ eV^2, $m_1 = 8.6 \times 10^{-3}$ eV, $y_0 = 0.02$, $r = 0.64$, $(\beta_\nu, \gamma_\nu) = (-100^\circ, -36^\circ)$ and find the current baryon asymmetry in the universe can be successfully explained. The numerical results are shown in Fig. 4. Here the relation $Y_B = 28/79\, Y_{B-L}$ [16] has been adopted. During the calculations, the parameters $\widehat{A}$, $\widehat{B}$ and $\widehat{C}$, which characterize the contribution of the triplet Higgs to the effective neutrino mass matrix (11), can also be determined by using Eq. (12): $\widehat{A} \simeq 0.41$, $\widehat{B} \simeq -0.093$ and $\widehat{C} \simeq 0.38$. In this case, it is easy to see that the right-handed neutrino provides a sizable contribution to the light neutrino masses.

In summary, we have proposed a minimal type-II seesaw model, which contains only one right-handed neutrino in addition to the triplet Higgs. Different from the usual type II seesaw models with several right-handed neutrinos, our model gives the CP asymmetry in the decay of the lightest right-handed neutrino just by the vertex correction involving the triplet Higgs, no matter whether the contribution of the triplet Higgs dominates the light neutrino masses. For illustration, the Fritzsch-type lepton mass matrices are considered. The neutrino oscillations and the baryon asymmetry can simultaneously be well explained. More phenomenological studies in this scenario are apparently interesting and desirable.

Acknowledgments

The authors would like to thank Wan-Lei Guo for useful discussions. This work is supported in part by the Ministry of Science and Technology of China under Grant No. NKBRSF G19990754 and by the National Natural Science Foundation of China. Gu would like to thank the hospitality of Department of Engineering Physics, Tsinghua University where part of this work was finished.

[1] For a review, see A. Strumia and F. Vissani, hep-ph/0606054.

[2] A. Strumia and F. Vissani, Nucl. Phys. B **726**, 294

(2005).
[3] D.N. Spergel *et al.*, astro-ph/0603449.
[4] M. Fukugita and T. Yanagida, Phys. Lett. B **174**, 45 (1986); P. Langacker, R.D Peccei, and T. Yanagida, Mod. Phys.Lett.A **1**, 541 (1986); M.A. Luty, Phys. Rev. D **45**, 455 (1992); R.N. Mohapatra and X. Zhang, Phys. Rev. D **46**, 5331 (1992).
[5] P. Minkowski, Phys. Lett. B **67**, 421 (1977); T. Yanagida, in *Proceedings of the Workshop on Unified Theory and the Baryon Number of the Universe*, edited by O. Sawada and A. Sugamoto (KEK, Tsukuba, 1979), p. 95; M. Gell-Mann, P. Ramond, and R. Slansky, in *Supergravity*, edited by F. van Nieuwenhuizen and D. Freedman (North Holland, Amsterdam, 1979), p. 315; S.L. Glashow, in *Quarks and Leptons*, edited by M. Lévy *et al.* (Plenum, New York, 1980), p. 707; R.N. Mohapatra and G. Senjanoviç, Phys. Rev. Lett. **44**, 912 (1980).
[6] J. Schechter and J.W.F. Valle, Phys. Rev. D **22**, 2227 (1980); Phys. Rev. D **25**, 774 (1982); G. Lazarides, Q. Shafi, and C. Wetterich, Nucl. Phys. B **181**, 287 (1981); R.N. Mohapatra and G. Senjanoviç, Phys. Rev. D **23**, 165 (1981); C. Wetterich, Nucl. Phys. B **187**, 343 (1981).
[7] E.J. Chun and S.K. Kang, Phys. Rev. D **63**, 097902 (2001); N. Sahu and U. Sarkar, hep-ph/0605007.
[8] T. Hambye and G. Senjanoviç, Phys. Lett. B **582**, 73 (2004).
[9] S. Antusch and S.F. King, Phys. Lett. B **597**, 199 (2004).
[10] H. Fritzsch, Phys. Lett. B **73**, 317 (1978); Nucl. Phys. B **155**, 189 (1979); Z.Z. Xing, Phys. Lett. B **550**, 178 (2002); M. Fukugita, M. Tanimoto and T. Yanagida, Phys. Lett. B **562**, 273 (2003); Z.Z. Xing and S. Zhou, Phys. Lett. B **593**, 156 (2004); S. Zhou and Z.Z. Xing, Eur. Phys. J. C **38**, 495 (2005).
[11] H. Nishiura, K. Matsuda and T. Fukuyama, Phys. Rev. D **60**, 013006 (1999); Z.Z. Xing and H. Zhang, Phys. Lett. B **569**, 30 (2003); J. Phys. G **30**, 129 (2004); and references therein.
[12] W.L. Guo, Phys. Rev. D **70**, 053009 (2004); W. Rodejohann, Phys. Rev. D **70**, 073010 (2004); N. Sahu and S. Uma Sankar, Phys. Rev. D **71**, 013006 (2004); W. Rodejohann and Z.Z. Xing, Phys. Lett. B **601**, 176 (2004).
[13] S. Eidelman *et al*, Phys. Lett. B **592**, 1 (2004).
[14] V.A. Kuzmin, V.A. Rubakov and M.E. Shaposhnikov, Phys. Lett. B **155**, 36 (1985); R.N. Mohapatra and X. Zhang, Phys. Rev. D **45**, 2699 (1992).
[15] P. Gu and X.J. Bi Phys. Rev. D **70**, 063511 (2004) and references therein.
[16] S.Yu. Khlebnikov and M.E. Shaposhnikov, Nucl. Phys. B **308**, 885 (1988).